

Remarks on some linear fractional stochastic equations

Ivan Nourdin

LPMA, Université Pierre et Marie Curie Paris 6,
Boîte courrier 188, 4 Place Jussieu, 75252 Paris Cedex 5, France
nourdin@ccr.jussieu.fr

Ciprian A. Tudor

SAMOS/MATISSE, Université de Panthéon-Sorbonne Paris 1
90, rue de Tolbiac, 75634 Paris Cédex 13, France
tudor@univ-paris1.fr

Abstract

Using the multiple stochastic integrals we prove an existence and uniqueness result for a linear stochastic equation driven by the fractional Brownian motion with any Hurst parameter. We study both the one parameter and two parameter cases. When the drift is zero, we show that in the one-parameter case the solution is an exponential, thus positive, function while in the two-parameter settings the solution is negative on a non-negligible set.

Key words: Fractional Brownian motion, fractional Brownian sheet, multiple stochastic integral, Girsanov transform.

2000 Mathematics Subject Classification: 60H05, 60G15, 60G18.

1 Introduction

The significant amount of applications where the fractional Brownian motion (fBm) is used led to the intensive development of the stochastic calculus with respect to this process and its planar version. The study of stochastic differential equations (SDEs) driven by a fractional Brownian motion followed in a natural way. Let us consider $(B_t^\alpha)_{t \in [0, T]}$ a fBm with Hurst parameter $\alpha \in (0, 1)$. Essentially, one can consider the SDE

$$dX_t = \sigma(t, X_t)dB_t^\alpha + b(t, X_t)dt \quad (1)$$

in two ways:

- the pathwise (Stratonovich) type (that is, the stochastic integral is considered in a pathwise sense);
- the divergence (Skorohod) type (that is, the stochastic integral is of divergence type).

The first type of equations, which includes the rough paths theory and the stochastic calculus via regularization, can in general be solved by now standard methods. We refer, among others, to [1, 4, 6, 8, 11, 12, 15, 19]. The second type (Skorohod stochastic equations) is more difficult to be solved. Even in the standard Brownian motion case (corresponding to $\alpha = 1/2$), we have an existence and uniqueness result only in two situations:

- when $\sigma(s, X_s) = \sigma(s)X_s$ with $\sigma(s)$ random: we then use an anticipating Girsanov transform, see [2],
- when $\sigma(s, X_s) = \sigma(s)X_s$ and $b(s, X_s) = b(s)X_s$ with σ, b two deterministic functions: we can then use a method based on the Wiener-Itô chaotic expansion.

This second approach will be considered in our paper. We will consider the stochastic equation

$$X_t = 1 + \int_{[0,t]} aX_s \delta B_s^\alpha + \int_{[0,t]} bX_s ds \quad (2)$$

where a, b are real numbers and the stochastic integral is understood in the Skorohod sense. We first prove existence and uniqueness results in the one-parameter case (that is when $t \in [0, T]$) and in the two-parameter case (that is when $t \in [0, T]^2$ and with B^α replaced by a fractional Brownian sheet $W^{\alpha, \beta}$ with Hurst parameters α, β).

Of course, the fact that the above linear equation can be solved by using Wiener-Itô multiple integrals is not very surprising; it has already used in [16] for $\alpha > \frac{1}{2}$. Nevertheless, we have to check some new technical aspects like: the proof of the case $\alpha \in (0, \frac{1}{2})$ or the proof of the two-parameter case for any Hurst parameters α and β .

More surprising is the behavior of the solution of (2) when the drift b is zero: in the one-parameter case, the solution is an exponential, hence positive, function while in the two-parameter case the solution is negative on a non-negligible set. We also mention that, comparing to the standard case when the Hurst parameters are $\frac{1}{2}$, new techniques like fractional Girsanov theorem and estimations of fractional norms of the kernels appearing in the chaotic expression of the solution of (2), are here needed.

We organized our paper as follows. Section 2 contains some preliminaries on fractional Brownian motion and fractional Brownian sheet. In Section 3 we study the existence, the uniqueness and the properties of the solution of equation (2) in both one-parameter and two-parameter cases. Section 4 contains a technical proof.

2 Preliminaries

Consider $(B_t^\alpha)_{t \in [0, T]}$ a fractional Brownian motion (fBm) with Hurst parameter $\alpha \in (0, 1)$ and let us denote by R^α its covariance function

$$R^\alpha(s, u) = \frac{1}{2} (s^{2\alpha} + u^{2\alpha} - |s - u|^{2\alpha}) \quad (3)$$

for every $s, u \in [0, T]$. It is well-known that B^α admits the Wiener integral representation

$$B_t^\alpha = \int_0^t K^\alpha(t, s) dW_s$$

where W denotes a standard Wiener process and

$$K^\alpha(t, s) = d_\alpha (t - s)^{\alpha - \frac{1}{2}} + s^{\alpha - \frac{1}{2}} F_1 \left(\frac{t}{s} \right), \quad (4)$$

d_α being a constant and $F_1(z) = d_\alpha \left(\frac{1}{2} - \alpha \right) \int_0^{z-1} \theta^{\alpha - 3/2} (1 - (\theta + 1)^{\alpha - 1/2}) d\theta$.

By $\mathcal{H}(\alpha)$ we will denote the canonical Hilbert space associated to B^α . That is, $\mathcal{H}(\alpha)$ is defined as the closure of the linear space generated by the indicator functions $\{1_{[0, t]}, t \in [0, T]\}$ with respect to the scalar product

$$\langle 1_{[0, t]}, 1_{[0, u]} \rangle_{\mathcal{H}(\alpha)} = R^\alpha(t, u). \quad (5)$$

The structure of $\mathcal{H}(\alpha)$ depends on the values of the Hurst parameter α . Let us recall the following facts:

- if $\alpha \in (\frac{1}{2}, 1)$, then it follows from [17] that the elements of $\mathcal{H}(\alpha)$ may be not functions but distributions of negative order. Thus it is more convenient to work with subspaces of $\mathcal{H}(\alpha)$ that are sets of functions. A such space is the set $|\mathcal{H}(\alpha)|$ of measurable functions on $[0, T]$ such that

$$\int_0^T \int_0^T |f(u)| |f(v)| |u - v|^{2\alpha - 2} dudv < \infty$$

endowed with the scalar product

$$\langle f, g \rangle_{|\mathcal{H}(\alpha)|} = \alpha(2\alpha - 1) \int_0^T \int_0^T f(u) f(v) |u - v|^{2\alpha - 2} dudv. \quad (6)$$

We have actually the inclusions

$$L^2([0, T]) \subset L^{\frac{1}{\alpha}}([0, T]) \subset |\mathcal{H}(\alpha)| \subset \mathcal{H}(\alpha). \quad (7)$$

- if $\alpha \in (0, \frac{1}{2})$ then the Hilbert space $\mathcal{H}(\alpha)$ is a space of functions contained in $L^2([0, T])$. It contains the space of Hölder functions of order $\alpha - \varepsilon$ with $\varepsilon > 0$ and it can be characterized by

$$\mathcal{H}(\alpha) = (K^*)^{-1}(L^2([0, T])) \quad (8)$$

where the operator K^* is given by

$$(K^*\varphi)(s) = K^\alpha(T, s)\varphi(s) + \int_s^T (\varphi(r) - \varphi(s)) \frac{\partial K^\alpha}{\partial r}(r, s) dr. \quad (9)$$

A fBm being a Gaussian process, it is possible to construct multiple Wiener-Itô stochastic integrals with respect to it. We refer to [13] for general settings or to [16] for the adaptation to the fractional Brownian motion case. We only recall that the multiple integral of order n (denoted by I_n) is an isometry from $\mathcal{U}^{\otimes n}$ to $L^2(\Omega)$ where \mathcal{U} is the Hilbert space $|\mathcal{H}(\alpha)|$ if $\alpha \in (\frac{1}{2}, 1)$ and the Hilbert space $\mathcal{H}(\alpha)$ if $\alpha \in (0, \frac{1}{2})$.

We need to introduce the space D^{ch} of stochastic processes that can be expressed in terms of multiple stochastic integrals. That is, we denote by D^{ch} the set of processes $u \in L^2(\Omega; \mathcal{U})$ such that for every $t \in [0, T]$,

$$u_t = \sum_{n \geq 0} I_n(f_n(\cdot, t))$$

where $f_n \in \mathcal{U}^{\otimes n+1}$ is symmetric in the first n variables and

$$\sum_{n \geq 1} (n+1)! \|f_n\|_{\mathcal{U}^{\otimes n+1}}^2 < \infty. \quad (10)$$

It follows from [16] (for $\alpha > \frac{1}{2}$) or [9] (for $\alpha < \frac{1}{2}$) that if $u \in D^{ch}$ then u is Skorohod integrable with respect to the fBm B^α and in this case its Skorohod integral is

$$\delta(u) = \sum_{n \geq 0} I_{n+1}(\tilde{f}_n) \quad (11)$$

where \tilde{f}_n means the symmetrization of f_n with respect to $n+1$ variables. Actually, in the case $\alpha < \frac{1}{2}$ the expression (11) corresponds to the divergence integral in the extended sense.

Let us consider now the two-parameter case. Here, $W^{\alpha, \beta}$ is a fractional Brownian sheet with Hurst parameters $\alpha, \beta \in (0, 1)$. Recall that $W^{\alpha, \beta}$ is defined as a centered Gaussian process starting from $(0, 0)$ with the covariance function

$$\begin{aligned} E \left(W_{s,t}^{\alpha, \beta} W_{u,v}^{\alpha, \beta} \right) &= R^{\alpha, \beta}(s, t, u, v) \\ &:= \frac{1}{2} (s^{2\alpha} + u^{2\alpha} - |s - u|^{2\alpha}) \frac{1}{2} (t^{2\beta} + v^{2\beta} - |t - v|^{2\beta}) \end{aligned}$$

and it can be represented as

$$W_{s,t}^{\alpha,\beta} = \int_0^t \int_0^s K^\alpha(t,u) K^\beta(s,v) dW_{u,v}$$

where $(W_{u,v})_{u,v \in [0,T]}$ is a standard Brownian sheet and K^α is given by (4). Denote by

$$K^{\alpha,\beta}(t,s) = K^\alpha(t,u) K^\beta(s,v).$$

and let $\mathcal{H}^{(2)}(\alpha,\beta) := \mathcal{H}^{(2)}$ be the canonical Hilbert space of the fractional Brownian sheet $W^{\alpha,\beta}$. That is, $\mathcal{H}^{(2)}$ is defined as the closure of the set of indicator functions $\{1_{[0,t] \times [0,s]}, t, s \in [0, T]\}$ with respect to the scalar product

$$\langle 1_{[0,t] \times [0,s]}, 1_{[0,u] \times [0,v]} \rangle_{\mathcal{H}^{(2)}} = R^{\alpha,\beta}(s,t,u,v) \quad (12)$$

for every $t, s, u, v \in [0, T]$.

By the above considerations, we will have:

- if $\alpha, \beta \in (\frac{1}{2}, 1)$, the elements of $\mathcal{H}^{(2)}$ may be not functions but distributions. Thus it is more convenient to work with subspaces of $\mathcal{H}^{(2)}$ that are sets of functions. We have actually the inclusions

$$L^2([0, T]^2) \subset |\mathcal{H}|^{(2)} \subset \mathcal{H}^{(2)} \quad (13)$$

where

$$|\mathcal{H}|^{(2)} = |\mathcal{H}(\alpha)| \otimes |\mathcal{H}(\beta)|$$

and $|\mathcal{H}(\alpha)|$ is defined by (6).

- if $\alpha, \beta \in (0, \frac{1}{2})$ then the canonical space $\mathcal{H}^{(2)}$ is a space of functions that can be written as

$$\mathcal{H}^{(2)} = (K^{*,2})^{-1} (L^2([0, T])) \subset L^2([0, T]) \quad (14)$$

where $K^{*,2}$ is the product operator $K^* \otimes K^*$ and K^* is given by (9).

- if $\alpha \in (\frac{1}{2}, 1)$ and $\beta \in (0, \frac{1}{2})$, then $|\mathcal{H}|^{(2)}$ is not a space of functions and we will work with the subspace $|\mathcal{H}(\alpha)| \otimes \mathcal{H}(\beta)$.

Let us denote by \mathcal{V} the Hilbert space: $|\mathcal{H}|^{(2)}$ if $\alpha, \beta \in (\frac{1}{2}, 1)$; $\mathcal{H}^{(2)}$ if $\alpha, \beta \in (0, \frac{1}{2})$ and $|\mathcal{H}(\alpha)| \otimes \mathcal{H}(\beta)$ if $\alpha \in (\frac{1}{2}, 1)$ and $\beta \in (0, \frac{1}{2})$.

We can of course consider multiple stochastic integrals with respect to the Gaussian process $W^{\alpha,\beta}$. Here the multiple integral of order n , still denoted by I_n , will be a isometry from $\mathcal{V}^{\otimes n}$ to $L^2(\Omega)$.

3 Linear stochastic equations with fractional Brownian motion and fractional Brownian sheet

Let us consider the following stochastic integral equation

$$X_t = 1 + \int_0^t a X_s \delta B_s^\alpha + \int_0^t b X_s ds, \quad t \in [0, T], \quad (15)$$

where $a, b \in \mathbb{R}$ and the stochastic integral above is considered in the Skorohod sense. We will first prove the existence and the uniqueness of the solution of (15), in the space D^{ch} . For $\alpha > \frac{1}{2}$ this has been proved in [16].

Proposition 1 *The equation (15) admits an unique solution $X \in D^{ch}$ given by*

$$X_t = \sum_{n \geq 0} I_n(f_n(\cdot, t)) \quad (16)$$

where the kernels f_n are given by

$$f_0(t) = e^{bt}$$

and for every $n \geq 1$,

$$f_n(t_1, \dots, t_n, t) = \frac{a^n}{n!} e^{bt} 1_{[0, t]}^{\otimes n}(t_1, \dots, t_n). \quad (17)$$

Proof: The expression (17) of the kernels f_n follows from Proposition 3.40 of [16]. One can also compute it easily by the recurrence relation

$$f_0(t) = e^{bt}, \quad f_n(t_1, \dots, t_n, t) = a \tilde{f}_{n-1}(t_1, \dots, t_{n-1}, t_n) 1_{[0, t]}(t_n), \quad \forall n \geq 1. \quad (18)$$

We only then need to prove that $f_n \in |\mathcal{H}(\alpha)|^{\otimes n+1}$ (if $\alpha > \frac{1}{2}$) and $f_n \in \mathcal{H}(\alpha)^{\otimes n+1}$ (if $\alpha < \frac{1}{2}$) and that the sum (10) converges.

If $\alpha > \frac{1}{2}$, this follows easily from the inclusion (7), since

$$\|\tilde{f}_n\|_{|\mathcal{H}(\alpha)|^{\otimes n+1}} \leq \text{cst} \|\tilde{f}_n\|_{L^2([0, T]^{n+1})}$$

and we can reduce to the classical situation ($\alpha = \frac{1}{2}$) where the result is known.

If $\alpha < \frac{1}{2}$, then we need a new proof because the norm $\mathcal{H}(\alpha)$ is bigger than the norm L^2 . Let us show that the kernel f_n given by (17) (viewed as a function of $n+1$ variables t_1, \dots, t_n, t) belongs to the space $\mathcal{H}(\alpha)^{\otimes n+1}$. Here we can adapt an argument used in [18]. We will show that

$$K^{*, n+1} f_n \in L^2([0, T]^{n+1})$$

where $K^{*, n}$ is the n times tensor product of K^* . It holds, by applying first the operator K^* to the variables t_1, \dots, t_n and then to the variable t ,

$$K^{*, n+1} f_n = \frac{a^n}{n!} K^* e^{bt} \left(K^{*, n} (1_{[0, t]}^{\otimes n}) \right)$$

and therefore, since

$$\|K^{*,n}(1_{[0,t]}^{\otimes n})\|_{L^2([0,T]^n)}^2 = \|K^*(1_{[0,t]})\|_{L^2([0,T])}^{2n} = t^{2n\alpha}$$

we get

$$\begin{aligned} \|K^{*,n+1}f_n\|_{L^2([0,T]^{n+1})} &= \frac{|a|^n}{n!} \|K^*(e^{bt}t^{2n\alpha})\|_{L^2([0,T])} \\ &= \frac{|a|^n}{n!} \left\| \sum_{k \geq 0} \frac{|b|^k}{k!} K^*(t^{k+2\alpha n}) \right\|_{L^2([0,T])}. \end{aligned}$$

Since for every $k \geq 1$, the function $t^{k+2\alpha n}$ is Lipschitz, then we have, using (9)

$$\begin{aligned} \|K^*(t^{k+2\alpha n})\|_{L^2([0,T])} &\leq C(\alpha, T)(k + 2\alpha n) \left[\int_0^T K^\alpha(T, t)^2 t^{4n\alpha + 2k} dt \right. \\ &\quad \left. + \int_0^T \left(\int_t^T (r-t) \frac{\partial K^\alpha}{\partial r}(r, t) dr \right)^2 dt \right]^{1/2} \\ &\leq C(\alpha, T)(k + 2\alpha n) T^{k+\alpha(2n+1)}. \end{aligned}$$

This implies that

$$\|K^{*,n+1}f_n\|_{L^2([0,T]^{n+1})} \leq \text{cst} \frac{|a|^n}{n!} T^{\alpha(2n+1)}. \quad (19)$$

The function f_n being symmetric in the first n variables, we have

$$\tilde{f}_n(t_1, \dots, t_{m+1}) = \frac{1}{m+1} \sum_{i=0}^{m+1} f_n(t_1, \dots, t_{m+1}^i, \dots, t_m)$$

where t_{m+1}^i means that t_{m+1} is on the position i . Clearly the bound (19) holds for \tilde{f}_n . By the above estimate, it is not difficult to see that the sum (10) is convergent because

$$\sum_{n \geq 0} (n+1)! \|\tilde{f}_n\|_{\mathcal{H}(\alpha)^{\otimes n+1}}^2 \leq \text{cst} \sum_{n \geq 0} \frac{a^{2n}}{n!} T^{2\alpha(2n+1)} < \infty.$$

The uniqueness of the solution in D^{ch} is obvious because, if there are two solutions, then the kernels of the chaotic expansion verifies both the relation (18). \blacksquare

In the particular case when the drift b is zero, we have the following

Corollary 1 *The unique solution in D^{ch} of the equation*

$$X_t = 1 + \int_0^t a X_s \delta B_s^\alpha, \quad t \in [0, T] \quad (20)$$

is given by

$$X_t = \exp \left(a B_t^\alpha - \frac{a^2}{2} t^{2\alpha} \right). \quad (21)$$

Proof: This is of course a consequence of Proposition 1. But, to compare to the two-parameter case, we prefer to show how the formula (21) is obtained. Let, for every $t \in [0, T]$

$$X_t = \sum_{n \geq 0} I_n(f_n(\cdot, t))$$

be the chaotic expression of X . Equation (20) can be rewritten as

$$\sum_{n \geq 0} I_n(f_n(\cdot, t)) = 1 + a \sum_{n \geq 0} I_{n+1} \left(f_n(\cdot, \star) \widetilde{1_{[0,t]}(\star)} \right) \quad (22)$$

where \cdot represents n variables, \star denotes one variable and $f_n(\cdot, \star) \widetilde{1_{[0,t]}(\star)}$ denotes the symmetrization of the function $f_n(\cdot, \star) 1_{[0,t]}(\star)$ in $n+1$ variables.

By identifying the corresponding Wiener chaos, we easily get

$$f_0(t) = 1, \quad f_1(t_1, t) = a 1_{[0,t]}(t_1)$$

and

$$f_2(t_1, t_2, t) = \frac{a^2}{2} (1_{[0,t_1]}(t_2) 1_{[0,t]}(t_1) + 1_{[0,t_2]}(t_1) 1_{[0,t]}(t_2)) = \frac{a^2}{2} 1_{[0,t]}^{\otimes 2}(t_1, t_2).$$

By induction we will get for every $n \geq 1$

$$f_n(t_1, t_2, \dots, t_n) = \frac{a^n}{n!} \sum_{i=0}^n 1_{[0,t_i]}^{\otimes n-1}(\hat{t}_i) 1_{[0,t]}(t_i) = \frac{a^n}{n!} 1_{[0,t]}^{\otimes n}(t_1, \dots, t_n) \quad (23)$$

where by \hat{t}_i we denoted the vector (t_1, \dots, t_n) with t_i missing. Therefore, we can express the solution of (20) as

$$X_t = \sum_{n \geq 0} \frac{a^n}{n!} I_n \left(1_{[0,t]}^{\otimes n} \right) = \exp \left(a B_t^H - \frac{a^2}{2} \|1_{[0,t]}\|_{\mathcal{H}}^2 \right) \quad (24)$$

where for the last equality we refer e.g. to [5]. ■

Remark 1 In Skorohod setting, it is difficult, in general, to write an Euler's type scheme associated to the equation $X_t = x_0 + \int_0^t \sigma(X_s) \delta B_s^\alpha$, even if $\alpha \geq 1/2$. Indeed, by using the integration by parts for the Skorohod integral δ and the Malliavin derivative D (see [13]) $\delta(Fu) = F\delta(u) - \langle DF, u \rangle_{\mathcal{H}(\alpha)}$ and by assuming that we approximate $X_{(k+1)/n}$ by $X_{k/n} + \int_{k/n}^{(k+1)/n} \sigma(X_{k/n}) \delta B_s^\alpha$ (as in the case $\alpha = 1/2$), one obtains

$$\widehat{X}_{(k+1)/n}^{(n)} = \widehat{X}_{k/n}^{(n)} + \sigma(\widehat{X}_{k/n}^{(n)}) \left(B_{(k+1)/n}^\alpha - B_{k/n}^\alpha \right) - \sigma'(\widehat{X}_{k/n}^{(n)}) \langle D\widehat{X}_{k/n}^{(n)}, 1_{[k/n, (k+1)/n]} \rangle_{\mathcal{H}(\alpha)}.$$

The problem is that the quantity $D\widehat{X}_{k/n}^{(n)}$ appears and that it is difficult to compute it directly (without knowing the solution). Moreover, standard Euler scheme do not apply

here because the L^2 -norm of the Skorohod integral involves the first Malliavin derivative which involves the second Malliavin derivative etc. and we cannot have closable formulas. In the linear case, taking advantage from the fact that we know explicitly the solution, we can see what the correct Euler scheme should be. Indeed, since we have $DX_{k/n} = aX_{k/n}\mathbf{1}_{[0,k/n]}$ (see Corollary 1 above), a natural Euler's type scheme associated to (20) is

$$\widehat{X}_{(k+1)/n}^{(n)} = \widehat{X}_{k/n}^{(n)} + a\widehat{X}_{k/n}^{(n)} \left(B_{(k+1)/n}^\alpha - B_{k/n}^\alpha \right) - \frac{a^2}{2} \widehat{X}_{k/n}^{(n)} \left[\left(\frac{k+1}{n} \right)^{2H} - \left(\frac{k}{n} \right)^{2H} - \frac{1}{n^{2H}} \right].$$

In fact, it is not very difficult to prove (using the same method as in the proof of Proposition 6 in [10]) that $(\widehat{X}_1^{(n)})$ converges in $L^2(\Omega)$ if and only if $\alpha \geq 1/2$ and that, in the case where $\alpha > 1/2$, the limit is $\exp\left(aB_1^\alpha - \frac{a^2}{2}\right)$.

As we have seen, the solution of (20) is an exponential, hence positive, function. We will show that the situation is different in the two-parameter case.

Before that, let us consider the equation corresponding to (15) in the two-parameter case

$$X_z = 1 + \int_{[0,z]} aX_r \delta W_r^{\alpha,\beta} + \int_{[0,z]} bX_r dr \quad (25)$$

where $z = (s, t) \in [0, T]^2$ and $W^{\alpha,\beta}$ is a fractional Brownian sheet with Hurst parameters $\alpha, \beta \in (0, 1)$.

We will denote now by $D^{ch,2}$ the class of functionals that can be represented as a serie of multiple stochastic integrals with respect to $W^{\alpha,\beta}$ (that is, $D^{ch,2}$ is the two-parameter equivalent of D^{ch}). In the next proposition, we show that (25) admits a unique solution in this space:

Proposition 2 *Let us denote by \mathcal{A}_n the set $\{(z_1, \dots, z_n) \in (\mathbb{R}^2)^n : \exists \sigma \in \mathfrak{S}_n, z_{\sigma(1)} \leq \dots \leq z_{\sigma(n)}\}$ (in the one-parameter case: $\mathcal{A}_n = \mathbb{R}^n$). If $z \in \mathcal{A}_n$, we consider $\sigma = \sigma_z \in \mathfrak{S}_n$ such that $z_{\sigma(1)} \leq \dots \leq z_{\sigma(n)}$.*

The equation (25) admits an unique solution $X \in D^{ch,2}$ given by $X_z = \sum_{n \geq 0} I_n(f_n(\cdot, z))$ where

$$f_n(z_1, \dots, z_n, z) = \frac{a^n}{n!} h_0(b(s - s_{\sigma_z(n)})(t - t_{\sigma_z(n)})) \times \mathbf{1}_{\mathcal{A}_n}(z_1, \dots, z_n) \mathbf{1}_{[0,z]}^{\otimes n}(z_1, \dots, z_n) \times \prod_{1 \leq j \leq n} h_0(b(s_{\sigma_z(j)} - s_{\sigma_z(j-1)})(t_{\sigma_z(j)} - t_{\sigma_z(j-1)})) \quad (26)$$

with $z = (s, t)$, $z_i = (s_i, t_i)$ and $h_0(x) = \sum_{n=0}^{\infty} \frac{x^n}{(n!)^2}$. We also used the convention that $\sigma_z(0) = 0$ and $z_0 = (0, 0)$.

Proof: We only prove the algebraic part (26) of the Proposition. Indeed, the fact that the kernels f_n belongs to $\mathcal{V}^{\otimes n+1}$ did not present new difficulties with respect to the proofs

of Propositions 1 and 3. Thus, we return to these proofs for this point. Let us write

$$X_z = \sum_{n \geq 0} I_n (f_n(\cdot, z)).$$

Here, I_n is the n -order Wiener-Itô multiple integral with respect to the fractional Brownian sheet $W^{\alpha, \beta}$ and $f_n \in L^2([0, T]^{2n})$. From (25) we have that $f_0(z) = h_0(bst)$ and for $n \geq 1$,

$$f_n(z_1, \dots, z_n, z) = a f_{n-1}(z_1, \dots, z_n) \widetilde{\mathbf{1}}_{[0, z]}(z_n) + b \int_{[0, z]} f_n(z_1, \dots, z_n, r) dr.$$

Let $n = 1$. We therefore have

$$f_1(z_1, z) = a h_0(b s_1 t_1) \mathbf{1}_{[0, z]}(z_1) + b \int_{[0, z]} f_1(z_1, r) dr$$

and

$$f_1(z_1, z) = a h_0(b s_1 t_1) h_0(b(s - s_1)(t - t_1)) \mathbf{1}_{[0, z]}(z_1)$$

hence (26) is satisfied. If $n = 2$ it holds that

$$\begin{aligned} f_2(z_1, z_2, z_3) \widetilde{\mathbf{1}}_{[0, z]}(z_3) &= \frac{1}{2} (a h_0(b s_1 t_1) h_0(b(s_2 - s_1)(t_2 - t_1)) \mathbf{1}_{0 \leq z_1 \leq z_2 \leq z} \\ &\quad + a h_0(b s_2 t_2) h_0(b(s_1 - s_2)(t_1 - t_2)) \mathbf{1}_{0 \leq z_2 \leq z_1 \leq z}). \end{aligned}$$

Since

$$f_2(z_1, z_2, z) = a f_1(z_1, z_2) \widetilde{\mathbf{1}}_{[0, z]}(z_2) + b \int_{[0, z]} f_1(z_1, r) dr$$

we deduce that

$$\begin{aligned} f_2(z_1, z_2, z) &= \frac{a^2}{2} (h_0(b s_1 t_1) h_0(b(s_2 - s_1)(t_2 - t_1)) h_0(b(s - s_2)(t - t_2)) \mathbf{1}_{0 \leq z_1 \leq z_2 \leq z} \\ &\quad + h_0(b s_2 t_2) h_0(b(s_1 - s_2)(t_1 - t_2)) h_0(b(s - s_1)(t - t_1)) \mathbf{1}_{0 \leq z_2 \leq z_1 \leq z}) \end{aligned}$$

and again (26) is verified. The above computations can be easily extended to an induction argument. \blacksquare

Let us now discuss the case $b = 0$:

Proposition 3 *The equation*

$$X_z = 1 + \int_{[0, z]} a X_r \delta W_r^{\alpha, \beta}, \quad z \in [0, T]^2 \quad (27)$$

admits an unique solution $X \in D^{ch, 2}$ given by $X_z = \sum_{n \geq 0} I_n(f_n(\cdot, z))$ where

$$f_n(\rho_1, \dots, \rho_n, z) = \frac{a^n}{n!} \sum_{i=1}^n \mathbf{1}_{[0, \rho_i]}^{\otimes n-1}(\hat{\rho}_i) \mathbf{1}_{[0, z]}(\rho_i). \quad (28)$$

Proof: Let us write

$$X_z = \sum_{n \geq 0} I_n (f_n(\cdot, z)).$$

From the equivalent of relation (22) in the two-parameter case, we obtain

$$f_0(z) = 1, \quad f_1(\rho_1, z) = a 1_{[0,z]}(\rho_1)$$

and in general relation (28) holds. Since $\mathcal{A}_n \neq (\mathbb{R}^2)^n$ (recall that \mathcal{A}_n is defined in Proposition 2), note that this last expression is not equal to $\frac{a^m}{m!} 1_{[0,z]}^{\otimes m}(\rho_1, \dots, \rho_n)$ as in the one-parameter case (see Corollary 1).

Let us now prove that the kernel f_n belongs to the space $\mathcal{V}^{\otimes n+1}$. When the Hurst parameters α and β are bigger than $\frac{1}{2}$, then we can use (13) and then refer to the standard case of the Brownian sheet. We will thus only discuss the case $\alpha, \beta < \frac{1}{2}$; the case $\alpha > \frac{1}{2}$ and $\beta < \frac{1}{2}$ will be a mixture of the other two cases. We use the induction. We will illustrate first the case $n = 2$. We check that $1_{[0,z]}(z_2)1_{[0,z_2]}(z_1)$ belongs to $\mathcal{H}^{(2)}^{\otimes 3}$. This actually reduces to proving that

$$1_{[0,t]}(t_2)1_{[0,t_2]}(t_1) \in \mathcal{H}(\alpha)^{\otimes 3}.$$

Let us apply the operator $K^{*,3}$ in three steps: first to the variable t_1 , then to the variable t and then to t_2 . It holds that

$$\begin{aligned} \|K^{*,3}(1_{[0,t]}(t_2)1_{[0,t_2]}(t_1))\|_{L^2([0,T]^3)}^2 &= \|K^{*,2}(t_2^{2\alpha}1_{[0,t]}(t_2))\|_{L^2([0,T]^2)}^2 \\ &= \|K^{*,1}(t_2^{2\alpha}(T-t_2)^{2\alpha})\|_{L^2([0,T])}^2 \end{aligned}$$

and to conclude we refer to Proposition 3.6 in [3]: it is a straightforward consequence of Lemma 4.3 in [3] that $(T-t_2)^{2\alpha}(B^\alpha)^2$ belongs to the extended domain of the divergence and therefore its expectation is in $\mathcal{H}(\alpha)$.

We will show now that the kernel $1_{[0,\rho_i]}^{\otimes n-1}(\hat{\rho}_i)1_{[0,z]}(\rho_i)$ has a finite norm in $\mathcal{H}^{(2)}^{\otimes n+1}$ by assuming that the result is true for n variables. It suffices to check that the function of $n+1$ (real) variables

$$1_{[0,t]}(t_n)1_{[0,t_n]}(t_{n+1}) \dots 1_{[0,t_2]}(t_1)$$

belongs to $\mathcal{H}(\alpha)^{\otimes n+1}$ or, equivalently, the operator $K^{*,n+1}$ applied to the above function is in $L^2([0,T]^{n+1})$. By applying first the operator K^* to the variable t_1 it holds that

$$\begin{aligned} &\|K^{*,n+1}(1_{[0,t]}(t_n)1_{[0,t_n]}(t_{n+1}) \dots 1_{[0,t_2]}(t_1))\|_{L^2([0,T]^{n+1})}^2 \\ &= \|K^{*,n}(1_{[0,t]}(t_n)1_{[0,t_n]}(t_{n+1}) \dots 1_{[0,t_3]}(t_2)t_2^{2\alpha})\|_{L^2([0,T]^n)}^2 \\ &= \|K^*(t_2^{2\alpha}g(t_2))\|_{L^2([0,T])}^2 \end{aligned}$$

where the function $t_2 \rightarrow g(t_2) := \|K^{*,n-1}(1_{[0,t]}(t_n)1_{[0,t_n]}(t_{n+1}) \dots 1_{[0,t_3]}(t_2))\|_{L^2([0,T]^{n-1})}^2$ belongs to $\mathcal{H}(\alpha)$ by the induction hypothesis. Now, we refer to the proof of Proposition 3.6 in [3] for the fact that $g(\cdot)E(B^\cdot)^2$ has a finite norm in $\mathcal{H}(\alpha)$.

It can actually be proved as above that

$$\|1_{[0,\rho_i]}^{\otimes n-1}(\hat{\rho}_i)1_{[0,z]}(\rho_i)\|_{\mathcal{H}^{(2)} \otimes n+1} \leq \frac{C^n}{n!}$$

for every n where C is a positive constant. Now we can finish as in proof of Proposition 1. \blacksquare

We will need the following Girsanov theorem. Its proof will be given in the Appendix.

Lemma 1 *For any $\varepsilon > 0$, the process*

$$W_{s,t}^{\alpha,\beta,\varepsilon} = W_{s,t}^{\alpha,\beta} - \frac{st}{\varepsilon} \quad (29)$$

has the same law as a fractional Brownian sheet with parameters α, β under the new probability P_ε given by

$$\frac{dP^\varepsilon}{dP} = \exp\left(\frac{1}{\varepsilon}W_{T,T}^{\alpha,\beta} - \frac{1}{2\varepsilon^2} \int_{[0,T]^2} \left(K_{\alpha,\beta}^{-1}(F(\cdot))(\rho)\right)^2 d\rho\right) \quad (30)$$

where $F(t, s) = ts$ and $K_{\alpha,\beta}$ is the operator associated to the kernel of the $W^{\alpha,\beta}$.

The solution of the equation (27) has actually a different behavior comparing to the one-parameter case (Corollary 1). We prove actually below that the solution of (27) is almost surely negative on a non-negligible set. Note that the same problem has been studied in the case of the standard Brownian sheet in [14].

Proposition 4 *Let X be the unique solution to (27) in the space $D^{ch,2}$. Then there exists an open set $\Delta \subset [0, T]^2$ such that*

$$P\{X_z < 0 \text{ for all } z \in \Delta\} > 0. \quad (31)$$

Proof: Note that the deterministic equation

$$g(s, t) = 1 + \int_0^s \int_0^t ag(u, v) dudv \quad (32)$$

admits the unique solution $g(s, t) = h_0(ast)$ with $h_0(x) = \sum_{n \geq 0} \frac{x^n}{(n!)^2}$ and that the function h_0 satisfy the property: there exists an open set $I = (-\beta, -\alpha)$ such that $h_0(x) < -\delta < 0$ for any $x \in I$ (see [14], page 231).

Suppose $a > 0$, fix $N > 0$ and define the open set

$$\Delta = \{(s, t), \alpha < ast < \beta, 0 < s, t < N\}.$$

For every $\varepsilon > 0$, consider

$$X_z^\varepsilon = 1 + \int_{[0,z]} a\varepsilon X_r^\varepsilon \delta W_r^{\alpha,\beta}.$$

Thanks to Corollary 1, we know that the solution X^ε of (3) is given by

$$X_z^\varepsilon = \sum_{n \geq 0} \varepsilon^n I_n(f_n(\cdot, z))$$

where the kernels f_n are given by $f_n(\rho_1, \dots, \rho_n, z) = \frac{a^n}{n!} \sum_{i=1}^n 1_{[0,\rho_i]}^{\otimes n-1}(\hat{\rho}_i) 1_{[0,z]}(\rho_i)$. Let us consider the equation

$$Y_z^\varepsilon = 1 + \int_{[0,z]} a\varepsilon Y_r^\varepsilon dW_r^{\alpha,\beta,\varepsilon} = 1 + \int_{[0,z]} a\varepsilon Y_r^\varepsilon dW_r^{\alpha,\beta} - \int_{[0,z]} aY_r^\varepsilon dr \quad (33)$$

and recall that, by Lemma 1, $W^{\alpha,\beta,\varepsilon}$ is a fractional Brownian sheet under P_ε . Now, we observe that

$$K = \sup_{\varepsilon > 0} \sup_z E |Y_z^\varepsilon|^2 < \infty. \quad (34)$$

In fact, to show that (34) holds is not difficult because it follows from Proposition 2 that the kernel of order n appearing in the chaotic expression of the solution of (33) are of the form ε^n multiplied to the kernel of order n of the solution of (33) with $\varepsilon = 1$. Then, $K \leq \sup_z E |Y_z^1|^2 < \infty$.

Now, by (32) and (33) we have, if $z = (t, s)$,

$$Y_{s,t}^\varepsilon - h_0(-ast) = -a \int_0^s \int_0^t (Y_{u,v}^\varepsilon - h_0(-auv)) dvdu + a\varepsilon \int_0^s \int_0^t Y_{u,v}^\varepsilon dW_{u,v}^{\alpha,\beta}$$

and using the bound (34) and the Gronwall Lemma in the plane we obtain

$$E \left[\sup_{s,t \in [0,T]} |Y_{s,t}^\varepsilon - h_0(-ast)|^2 \right] \rightarrow_{\varepsilon \rightarrow 0} 0.$$

Since $h_0(-ast) < -\delta$ for $(s, t) \in \Delta$, it follows that

$$P(Y_z^\varepsilon < 0, \forall z \in \Delta) \rightarrow_{\varepsilon \rightarrow 0} 1.$$

Thus, for every $\varepsilon > 0$ small enough

$$P(Y_z^\varepsilon < 0, \forall z \in \Delta) > 0$$

and

$$P_\varepsilon(Y_z^\varepsilon < 0, \forall z \in \Delta) = P(X_z^\varepsilon < 0, \forall z \in \Delta) > 0.$$

Since $W_{c_1 s, c_2 t}^{\alpha,\beta}$ has the same law as $c_1^\alpha c_2^\beta W_{s,t}^{\alpha,\beta}$ as process, we get that $X_{s,t}^\varepsilon$ is equal in law to $X_{\varepsilon^{2\alpha} s, \varepsilon^{2\beta} t}$. So, for $\varepsilon > 0$ small enough,

$$P(X_{\varepsilon^{2\alpha} s, \varepsilon^{2\beta} t} < 0, \forall z \in \Delta) > 0$$

and the conclusion follows. ■

4 Appendix

Proof of Lemma 1: The conclusion will follow from the Girsanov theorem for the fractional Brownian sheet (see Theorem 3 in [7]) if we show that the functions $F(s, t) = st$ belongs to the space $I^{\alpha+\frac{1}{2}, \beta+\frac{1}{2}}(L^2([0, T]^2))$ or equivalently,

$$K_{\alpha, \beta}^{-1}(F(\cdot)) \in L^2([0, T]^2).$$

To show this, we will need the expression of its inverse operator in terms of fractional integrals and derivatives (see e.g. [7])

$$K_{\alpha, \beta}^{-1}h(t, s) = t^{\alpha-\frac{1}{2}}s^{\beta-\frac{1}{2}}I^{\frac{1}{2}-\alpha, \frac{1}{2}-\beta} \left(t^{\frac{1}{2}-\alpha}s^{\frac{1}{2}-\beta} \frac{\partial^2 h}{\partial s \partial t} \right), \quad \alpha, \beta < \frac{1}{2} \quad (35)$$

and

$$K_{\alpha, \beta}^{-1}h(t, s) = t^{\alpha-\frac{1}{2}}s^{\beta-\frac{1}{2}}D^{\alpha-\frac{1}{2}, \beta-\frac{1}{2}} \left(t^{\frac{1}{2}-\alpha}s^{\frac{1}{2}-\beta} \frac{\partial^2 h}{\partial s \partial t} \right), \quad \alpha, \beta > \frac{1}{2}. \quad (36)$$

Here,

$$I^{\alpha, \beta}f(x, y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^x \int_0^y (x-u)^{\alpha-1}(y-v)^{\beta-1}f(u, v)dudv$$

and

$$D^{\alpha, \beta}f(x, y) = \frac{1}{\Gamma(1-\alpha)\Gamma(1-\beta)} \frac{\partial^2}{\partial x \partial y} \int_0^x \int_0^y \frac{f(u, v)}{(x-u)^\alpha(y-v)^\beta} dudv,$$

with Γ the Euler function.

For $\alpha, \beta \in (0, \frac{1}{2})$ we have

$$\begin{aligned} K_{\alpha, \beta}^{-1}F(t, s) &= t^{\alpha-\frac{1}{2}}s^{\beta-\frac{1}{2}}I^{\alpha-\frac{1}{2}, \beta-\frac{1}{2}} \left(t^{\frac{1}{2}-\alpha}s^{\frac{1}{2}-\beta} \right) \\ &= \frac{1}{\Gamma(\frac{1}{2}-\alpha)\Gamma(\frac{1}{2}-\beta)} t^{\alpha-\frac{1}{2}}s^{\beta-\frac{1}{2}} \int_0^t \int_0^s (t-u)^{-\frac{1}{2}-\alpha}u^{\frac{1}{2}-\alpha}(s-v)^{-\frac{1}{2}-\beta}v^{\frac{1}{2}-\beta}dvdu \end{aligned}$$

and this belongs to $L^2([0, T]^2)$.

If $\alpha, \beta \in (\frac{1}{2}, 1)$ then by (36) we can write

$$\begin{aligned}
& K_{\alpha, \beta}^{-1} F(t, s) \\
= & t^{\alpha - \frac{1}{2}} s^{\beta - \frac{1}{2}} D^{\alpha - \frac{1}{2}, \beta - \frac{1}{2}} \left(t^{\frac{1}{2} - \alpha} s^{\frac{1}{2} - \beta} \right) (t, s) \\
= & t^{\alpha - \frac{1}{2}} s^{\beta - \frac{1}{2}} \frac{1}{\Gamma(\frac{3}{2} - \alpha) \Gamma(\frac{3}{2} - \beta)} \left[\frac{t^{\frac{1}{2} - \alpha} s^{\frac{1}{2} - \beta}}{t^{\alpha - \frac{1}{2}} s^{\beta - \frac{1}{2}}} \right. \\
& + \frac{\alpha - \frac{1}{2}}{s^{\beta - \frac{1}{2}}} \int_0^t \frac{t^{\frac{1}{2} - \alpha} s^{\frac{1}{2} - \beta} - u^{\frac{1}{2} - \alpha} s^{\frac{1}{2} - \beta}}{(t - u)^{\alpha + \frac{1}{2}}} du \\
& + \frac{\beta - \frac{1}{2}}{t^{\alpha - \frac{1}{2}}} \int_0^s \frac{t^{\frac{1}{2} - \alpha} s^{\frac{1}{2} - \beta} - t^{\frac{1}{2} - \alpha} v^{\frac{1}{2} - \beta}}{(s - v)^{\beta + \frac{1}{2}}} dv \\
& \left. + (\alpha - \frac{1}{2})(\beta - \frac{1}{2}) \int_0^t \int_0^s \frac{t^{\frac{1}{2} - \alpha} s^{\frac{1}{2} - \beta} - u^{\frac{1}{2} - \alpha} s^{\frac{1}{2} - \beta} - t^{\frac{1}{2} - \alpha} v^{\frac{1}{2} - \beta} + u^{\frac{1}{2} - \alpha} v^{\frac{1}{2} - \beta}}{(t - u)^{\alpha + \frac{1}{2}} (s - v)^{\beta + \frac{1}{2}}} dv du \right].
\end{aligned}$$

Since

$$\int_0^t \frac{t^{\frac{1}{2} - \alpha} - u^{\frac{1}{2} - \alpha}}{(t - u)^{\alpha + \frac{1}{2}}} du = c(\alpha) t^{1 - 2\alpha},$$

it is not difficult to see that the above function is in $L^2([0, T]^2)$.

If $\alpha \in (0, \frac{1}{2})$ and $\beta \in (\frac{1}{2}, 1)$, then we have

$$\begin{aligned}
K_{\alpha, \beta}^{-1} F(t, s) &= C(\alpha, \beta) t^{\alpha - \frac{1}{2}} \int_0^t (t - u)^{-\frac{1}{2} - \alpha} u^{\frac{1}{2} - \alpha} du \\
&\quad \times s^{\frac{1}{2} - \beta} + (\beta - \frac{1}{2}) \int_0^s \frac{t^{\frac{1}{2} - \alpha} s^{\frac{1}{2} - \beta} - t^{\frac{1}{2} - \alpha} v^{\frac{1}{2} - \beta}}{(s - v)^{\beta + \frac{1}{2}}} dv
\end{aligned}$$

and the conclusion is clearly a consequence of the above two cases. The proof of Lemma is done. \blacksquare

References

- [1] E. Alos, J.A. Leon and D. Nualart (2001): *Stratonovich calculus for fractional Brownian motion with Hurst parameter less than $\frac{1}{2}$* . Taiwanese Journal of Math., 4, pag. 609-632.
- [2] R. Buckdahn (1991): *Linear stochastic Skorohod differential equations*. Probab. Theory Rel. Fields, 90, pag. 223-240.
- [3] P. Cheridito and D. Nualart (2005): *Stochastic integration of divergence type with respect to the fractional Brownian motion with Hurst parameter $H \in (0, \frac{1}{2})$* . Ann. Inst. H. Poincaré Probab. Statist., 41 (6), pag. 1049-1081.

- [4] L. Coutin and Z. Qian (2002): *Stochastic analysis, rough path analysis and fractional Brownian motion*. Prob. Theory Rel. Fields, 122(1), pag. 108-140.
- [5] T. E. Duncan, Y. Hu and B. Pasik-Duncan (2000): *Stochastic calculus for fractional Brownian motion I. Theory*. Siam J. Control Optim., 38(2), pag. 582-612.
- [6] M. Errami, F. Russo (2003). *n-covariation and symmetric SDEs driven by finite cubic variation process*. Stoch. Processes and their Applications, 104, pag. 259-299.
- [7] M. Erraoui, D. Nualart and Y. Ouknine (2003): *Hyperbolic stochastic partial differential equations with additive fractional Brownian sheet*. Stochastic and Dynamics, 3, pag. 121-139.
- [8] D. Feyel, A. De La Pradelle (2003): *Curvilinear integrals along enriched paths*. Preprint Université d'Évry.
- [9] J.A. Leon and D. Nualart (2005): *An extension of the divergence operator for Gaussian processes*. Stoch. Processes and their Applications, 115, pag. 481-492.
- [10] I. Nourdin (2005): *Schémas d'approximation associés à une équation différentielle dirigée par une fonction höldérienne; cas du mouvement brownien fractionnaire*. C.R. Math. Acad. Sci. Paris, Ser. I, 340, pag. 611-614.
- [11] I. Nourdin (2005): *A simple theory for the study of SDEs driven by a fractional Brownian motion, in dimension one*. Preprint LPMA, Paris 6.
- [12] I. Nourdin and T. Simon (2005): *Correcting symmetric integrals by a Lévy area*. Preprint LPMA, Paris 6.
- [13] D. Nualart (1995): *The Malliavin calculus and related topics*. Springer.
- [14] D. Nualart (1987): *Some remarks on a linear stochastic differential equation*. Stat. and Probab. Letters, 5, pag. 231-234.
- [15] D. Nualart and A. Răşcanu (2002). *Differential equations driven by fractional Brownian motion*. Collect. Math., 53 (1), pag. 55-81.
- [16] V. Perez Abreu and C. Tudor (2002): *A transfer principle for multiple stochastic fractional integrals*. Bol. Soc. Mat. Mexicana, 8(3), pag. 187-203.
- [17] V. Pipiras and M. Taqqu (2000): *Integration questions related to fractional Brownian motion*. Probab. Theory Rel. Fields, 118, pag. 251-291.
- [18] C.A. Tudor and F. Viens (2004): *Itô formula for the fractional Brownian sheet using the extended divergence operator*. Preprint SAMOS Paris 1.
- [19] M. Zaehle (1998): *Integration with respect to fractal functions and stochastic calculus*. Prob. Theory Rel. Fields, 111, pag. 333-374.